



Research Paper

*Afr. J. Traditional,
Complementary and Alternative
Medicines*
www.africanethnomedicines.net

ISSN 0189-6016©2009

PHYTOCHEMICAL AND ANTIMICROBIAL STUDIES OF FOUR SPECIES OF *COLA* SCHOTT & ENDL. (STERCULIACEAE)Mubo Adeola Sonibare¹, Micheal O. Soladoye², Oyedokun O. Esan², Oluwadayo O. Sonibare³¹Department of Pharmacognosy, Faculty of Pharmacy, University of Ibadan, Nigeria²Department of Plant Science and Applied Zoology, Olabisi Onabanjo University, Ago-Iwoye, Nigeria³Department of Chemistry, University of Ibadan, Nigeria*E-mail: sonibaredeola@yahoo.com

Abstract

The *in-vitro* antimicrobial evaluation of ethanol extracts of four species of *Cola* Schott & Endl. was done using human isolated strains of *Staphylococcus aureus*, *Staphylococcus albus*, *Bacillus subtilis*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa*, *Candida albicans*, *Aspergillus niger* as test organisms. The assays were carried out by agar well diffusion, erythromycin and ketoconazole served as the control drugs. The leaf ethanol extracts of the plants were found to be more effective against the tested fungi than the bacteria at high concentrations. None of the extracts was active against *Staphylococcus aureus*. Plant extract of *C. acuminata* (P. Beauv.) Schott & Endl. and *C. nitida* (Vent) Schott & Endl. showed activity on *S. albus* at concentrations ranging from 10-150 mgml⁻¹ having comparable diameters of zone of inhibition of 7.3±0.03-16.0±0.0 for *C. acuminata* and 10.0±0.0-19.0±0.0 for *C. nitida*. Also, these two species of *Cola* demonstrated activities on *C. albicans* and *A. niger* at concentrations ranging from 90-150mgml⁻¹ with relatively close diameters of zone of inhibition. Only *C. acuminata* inhibited the growth of *K. pneumoniae* at the MIC of 90mgml⁻¹ whereas, *C. albicans* was inhibited by *C. acuminata*, *C. millenii* K. Schum and *C. gigantea* A.Chev. at the MIC of 120mgml⁻¹. Phytochemical screening of the four species of *Cola* showed the presence of alkaloids, saponins, tannins and cardenolides in all the plants which apart from showing the probable closeness of the species could also be responsible for the observed activities. The antimicrobial property shown by the plant extracts is an evidence of the ethnomedicinal uses of the plants. The similarity observed in the phytochemical constituents and antimicrobial activities demonstrated by *C nitida* (Vent.) Schott & Endl., *C. millenii* and *C.gigantea* A. Chev. and *C. acuminata* suggest a probable closeness among these species. The results obtained in this study provide preliminary evidence of the chemotaxonomic significance of secondary metabolites and antimicrobial activities in infra-generic taxonomy of species of *Cola*.

Key words: *Cola* species, Sterculiaceae, phytochemical screening, antimicrobial studies, taxonomy.

Introduction

Cola Schott & Endl. (Sterculiaceae) is a genus of about 125 species of trees indigenous to the tropical rain-forest African region (Ratsch, 2005). Phylogenetic information reveals that the genus was formerly classified in the family Malvaceae, subfamily Sterculioideae and was later transferred into the separate family Sterculiaceae. *Cola* is one of the largest in the family Sterculiaceae and is related to the South American genus *Theobroma*. It comprises of evergreen moderately sized trees often growing to a height of 20m with glossy ovoid leaves up to 30cm long. *Cola* species are found mostly in the relatively dry parts of the rain forest, although *Cola millenii* and *Cola gigantea* are widely distributed in wet and dry forest environments (Kuoame and Sacande, 2006; Olorode, 1984).

According to Russell (1955), the systematics of *Cola* species was in a state of “indescribable confusion”. In an attempt to resolve this confusion, Chevalier and Perrot (1911) created the Subgenus *Eucola*

containing five species of edible kolanuts – *Cola nitida* (important for trade), *Cola acuminata* (important for socio-cultural values), *Cola ballayi*, *Cola verticillata* and *Cola sphaerocarpa*. The latter three species are not known to be cultivated

The mature fruit of *Cola* species is a nut known as kolanut (Duke, 2001). It has a bitter flavour and high caffeine content (Blades, 2000; Benjamin *et al.*, 1991). It is chewed in many West African cultures individually or in a group setting. It is often used ceremonially, presented to tribal chiefs or to guests. Chewing kolanut can ease hunger pangs. Kolanuts are used mainly for their stimulant and euphoriant qualities. They have effects similar to other xanthine containing herbs like cocoa, tea etc. However, the effects are distinctively different, producing a stronger state of euphoria and well being (Benjamin *et al.*, 1991). They have stimulant effects on the central nervous system and heart. Kolanuts are used as a source of alkaloids in pharmaceutical preparations (Newall *et al.*, 1996; Bradley, 1992; Opeke, 1992).

Various medicinal and pharmacological values have been observed in species of *Cola* (Daels-Rakotoarison *et al.*, 2003; Steinegger and Hansel, 1992). Kolanuts are often used to treat whooping cough and asthma. The caffeine present acts as a bronchodilator, expanding the bronchial air passages (Jayeola, 2001; Kim, 2001). Kolanuts are also employed in the treatment of malaria and fever (Odugbemi, 2006). Experiments using animals indicate that kolanuts have analeptic and lipolytic properties and stimulate the secretion of gastric juices (GRIN, 2007).

Odugbemi (2006) reported that the leaves of *Cola millenii* are used in the treatment of ringworm, scabies, gonorrhoea, dysentery and ophthalmia. Traditionally, the leaves, twigs, flowers, fruit follicles and the bark of *Cola nitida* and *Cola acuminata* are used to prepare a tonic as a remedy for dysentery, coughs, diarrhoea, vomiting and chest complaints (Burkill, 1995; Irvine, 1961).

This paper reports the phytochemical and antimicrobial activities of four species of *Cola* with a possible evaluation of their chemotaxonomic significance.

Materials and methods

Plant materials

Fresh leaves of *Cola acuminata* was collected in Ibadan, Oyo State, Nigeria. *Cola nitida*, *Cola millenii* and *Cola gigantea* were collected at different location in Ago-Iwoye, Ogun State, Nigeria in June 2007. The plants were identified and authenticated by Mr. T.K Odewo at the Forest Herbarium Ibadan (FHI) where the voucher specimens were also deposited under the following numbers: *Cola acuminata* (P. Beauv.) Schott & Endl. FHI 107892, *Cola milleni* K. Schum. FHI 107893, *Cola nitida* (Vent) Schott & Endl. FHI 107894, *Cola gigantea* A. Chev. var. *gigantea* Bull. FHI 107895. The voucher information which includes locality of collection and herbarium numbers of the *Cola* species are presented in Table 1.

Extraction of Plant Material

The air-dried and powdered leaves were extracted by maceration of 200g of the dried, pulverized leaves at room temperature for 48 hours in 2.5 litres of 96% ethanol. The mixture was filtered using Whatman No. 1 filter paper and the filtrate solution was evaporated in a water bath at 70°C to obtain a paste. The extracts gave yields of 3.9%, 0.6%, 2.8% and 2.5%, respectively, for *C. nitida*, *C. acuminata*, *C. millenii* and *C. gigantea*.

Phytochemical screening

The dried, pulverized leaves were subjected to phytochemical analysis to screen for the presence of secondary metabolites such as alkaloids, saponins, anthraquinones, cardenolides and tannins. The phytochemical screening was carried out using standard procedure (Ajaiyeoba *et al.*, 2003; Trease and Evans, 1989). Brief description is as follows:

Alkaloids: 70ml of 10% HCl was added to 4g of each sample in appropriately labeled conical flasks and boiled for 10 mins. Each boiled sample was filtered and allowed to cool. The filtrates were poured into four labeled test tubes. Few drops of Dragendoff's, Mayer's, Wagner's reagents were added to each test tube separately. Alkaloids were recorded as present in the sample if turbidity or a brownish precipitate was observed.

Saponins: 4g of each sample was dissolved in distilled water and heated for 2-5 mins. The mixtures were filtered, allowed to cool and shaken continuously for 2 mins to induce the production of froth. They were then left to stand for 15 mins. The observation of frothing was indicative of presence of saponin.

Test for Tannins: 1g of each sample was heated with 20ml of water for 5 mins in appropriately labeled test-tubes. Each solution was allowed to cool and then filtered. 1ml of each filtrate was diluted with 5ml distilled water in a test tube; few drops of 0.1% ferric chloride solution were added. A characteristic blue, blue-black, green or blue-green colour and precipitate indicate the presence of tannin.

Anthraquinones: 1g of each sample was shaken with 10ml of ferric chloride solution mixed with 5ml of HCL. Each mixture was heated in a water bath for 10-15 mins, filtered and allowed to cool. The filtrate was extracted with chloroform and shaken gently. The clear layers at the base were pipette into test tubes and 2ml each of ammonia solution was added. An observation of a delicate pink rose colour indicated the presence of anthraquinones.

Cardenolides: 4g of each sample were extracted in test tubes with 80% ethanol, all appropriately labeled. They were then divided into two portions for Kedde's test and Keller-Killiani's test. For Kedde's Test, few drops of 10% lead acetate were added to each of the tubes, followed by few drops of distilled water and chloroform. The contents were then evaporated to dryness in a water bath. 5% NaOH was added to each residue and then 2% of 3,5 dinitrobenzene acid. For Keller – Killiani's Test, few drops of 10% lead acetate, water and chloroform were added to each test sample. The mixtures were also evaporated to dryness in a water bath and subsequently, few drops of concentrated sulphuric acid were added. For Keller-Killiani's test, a brown ring indicated the presence of cardenolides while for the Kedde's test, a brown to purple colour was indicative of presence of cardenolides.

Antimicrobial screening

Microorganisms: Four human pathogenic bacteria made up of two Gram-positive (*Staphylococcus aureus*, *Staphylococcus albus*: ATCC: 27856 (Wilson and Stuart, 1965; Stich, 1932), and two Gram-negative (*Klebsiella pneumoniae*, *Pseudomonas aeruginosa*: ATCC: 27856) and *Bacillus subtilis* (ATCC: 1457), were used for the antibacterial assay. One yeast (*Candida albicans* (MTCC: 227) and one mold (*Aspergillus niger* (MTCC: 227) were used for the antifungal assay. All the organisms were local isolates from the Laboratory bacterial stock of the Department of Plant Science and Applied Zoology, Olabisi Onabanjo University, Ogun State, Nigeria. Three to five identical colonies from stored slopes of microorganisms (bacteria and fungi) were lifted with a sterile wire loop and transferred into a 5ml single strength nutrient broth (Biochemica, Spain) contained in well labeled screw cap bottles for each bacterium and fungus respectively. The bottles were well shaken and incubated at room temperature for 18-24h for bacteria and 72h for fungi.

The agar well diffusion method was used to test the plant extracts for antimicrobial activity. Briefly, 15ml of melted and cooled nutrient agar (Himedia Laboratories, India) and potato dextrose agar (Himedia Laboratories, India) were added to 0.2ml of 1 in 100 dilutions of bacteria and fungal cultures respectively in sterile Petri dishes. The contents were mixed. After the agar in each plate solidified, 6 wells of 5mm each were bored in each plate using an aseptic cork borer. 0.1ml of plant extracts at varying concentrations (10mgml⁻¹, 30mgml⁻¹, 90mgml⁻¹, 120mgml⁻¹, 150mgml⁻¹) as well as the standard antibiotic solution was loaded into the wells. Control experiments were set up using erythromycin and ketoconazole (2mgml⁻¹) for the bacterial and fungal assays respectively. The plates were incubated at 37^oC for 24h for bacteria and 48h for fungi. All inoculation procedures were undertaken under aseptic conditions. According to pharmacological and biometric specifications, the antimicrobial studies were done in triplicates. With the aid of a transparent ruler the diameters of zones of inhibition around the wells were measured in mm for all the three replicates and the average of the three measurements was calculated as an indication of activity.

The minimum inhibitory concentration (MIC) of plant extracts was determined using the broth dilution method as described by Sahm and Washington (1990). Briefly, 1ml of the extract solution at the concentration of 120mgml⁻¹ was added to 1ml of nutrient broth and subsequently transferred to make solutions of varying concentrations (120, 90, 60, 10 mgml⁻¹) in different test tubes. Then 1ml of bacterial and fungal suspensions and 0.1ml of plant extracts at the different concentrations was added to each test tube and incubated at 37^oC for 24h for bacteria and 48h for fungi. The test tube with the concentration of plant extract at which no detectable growth was observed was considered as the MIC.

Results

The result of the phytochemical screening of the leaf samples is presented in Table 2. The secondary metabolites tested for were alkaloids, saponins, tannins, cardenolides and anthraquinones. The result shows that alkaloids, saponins and tannins are present in all the four species. Anthraquinones are absent in all species except *C. gigantea*. Cardenolides are present in *C. acuminata*, *C. nitida* and *C. gigantea* but absent in *C. millenii*.

The results of the antimicrobial screening of the ethanol crude plant extracts of the species are presented in Table 3 while Table 4 shows the minimum inhibitory concentration (MIC) of each extract. The plant extracts were found to be more effective on the tested fungi than they were on bacteria. All the extracts showed important inhibition of fungal growth at the concentrations of 90, 120 and 150mgml⁻¹. Only a few bacteria were susceptible to the extracts at high concentration.

No antibacterial activity was observed against *Staphylococcus aureus*, a Gram-positive bacterium. Only the crude extract of *C. acuminata* was active against *Klebsiella pneumoniae*, (a Gram-negative bacterium) with

the diameter of the zone of inhibition of 9.6mm and the MIC of 90mgml⁻¹. Likewise, the extract of *C. millenii* was the only extract that inhibited the growth of *Pseudomonas aeruginosa* (a Gram negative bacterium) at a minimum diameter of zone of inhibition of 8.3mm and an MIC of 90mgml⁻¹. *Bacillus subtilis* was resistant to all the extracts, except to that of *C. gigantea*, which was active against the organism at an MIC of 10mgml⁻¹. Extracts of *C. acuminata* and *C. nitida* were active against the two fungi *Aspergillus niger* and *Candida albicans*, both showed activity at MIC of 90mgml⁻¹. *C. millenii* extract inhibited the growth of *Aspergillus niger* at an MIC 90mgml⁻¹ but was only able to inhibit the growth of *Candida albicans* at higher concentrations of 120mgml⁻¹ and 150mgml⁻¹. The extract of *C. gigantea* was active against *Aspergillus niger* and *Candida albicans* at MICs of 60mgml⁻¹ and 90mgml⁻¹ respectively.

Table 1: Voucher information of representatives of *Cola* species studied

Binomials and Authorities	Locality of collection	Herbarium Number
<i>Cola acuminata</i> (P.Beauv.) Schott & Endl.	Ibadan	FHI 107892
<i>Cola millenii</i> K.Schum.	Ago-Iwoye	FHI 107893
<i>Cola nitida</i> (Vent) Schott & Endl.	Ago-Iwoye	FHI 107894
<i>Cola gigantea</i> A.Chev. var <i>gigantea</i> Bull.	Ago-Iwoye	FHI 107895

Table 2. Phytochemical constituents of crude extract of *Cola* species

Secondary Metabolites	Plant Samples			
	<i>C. acuminata</i>	<i>C. nitida</i>	<i>C. millenii</i>	<i>C. gigantea</i>
Alkaloids	+	+	+	+
Saponins	+	+	+	+
Tannins	+	+	+	+
Cardenolides	+	+	-	+
Anthraquinones	-	-	-	+

+: Presence of secondary metabolite

-: Absence of secondary metabolite

Discussion

The presence of the secondary metabolites (alkaloids, saponins and tannins) in *Cola acuminata*, *Cola nitida*, *Cola millenii* and *Cola gigantea* partly enhances the chemotaxonomic characterization of the four *Cola* species. Although the presence of similar secondary metabolites may not necessarily justify the generic closeness of these species, it is a noteworthy observation that requires further studies in attempting to resolve the relationship among the three species. Owing to the presence of these three secondary metabolites and the similarity of their occurrence, it can be proposed that the four *Cola* species are appropriately classified into the

same genus. This suggests that the four plants are closely related and could be assumed to have a common origin, a claim supported by the previous work (Adegoke et al., 1968; Keay, 1989; Oliver-Bever, 1986; Purseglove, 1968; Oliver, 1960). This claim could be strengthened with a further evaluation of the active principles responsible for the antimicrobial activities observed in these species.

The environment is known to potentially influence the morphology and expression of compounds in plants (Folkers et al., 2008; Shen et al., 2008; Tsukaya et al., 2007; Braga et al., 2006; Cybulskill et al., 2000). Plant physiologists have reported that a particular compound may be produced only at certain times or under certain conditions. This may be the case with *Cola gigantea*, the only plant that showed a rare presence of anthraquinones. *C. gigantea* is mainly found in forests while the other three species are more domesticated in

Table 3: Antimicrobial activity of crude ethanol extracts of four *Cola* species Zones of Inhibition (mm)

Plant Sp	Yield (%)	Conc. mgml ⁻¹	<i>S. albus</i>	<i>S. aureus</i>	<i>B. subtilis</i>	<i>K. pneumoniae</i>	<i>P. aeruginosa</i>	<i>C. albicans</i>	<i>A. niger</i>
<i>C. acuminata</i>	0.6	10	10.0±0.0	-	-	-	-	-	-
		30	11.0±0.0	-	-	-	-	-	-
		60	7.3±0.03	-	-	-	-	-	-
		90	13.0±0.0	-	-	20.5±1.0	-	13.5±0.0	13.5±0.0
		120	12.5±1.0	-	-	18.5±1.0	-	15.5±1.0	15.5±1.0
		150	16.0±0.0	-	-	9.6±1.1	-	19.5±1.0	19.5±1.0
<i>C. nitida</i>	3.9	10	10.0±0.0	-	-	-	-	-	-
		30	13.0±0.0	-	-	-	-	-	-
		60	19.0±0.0	-	-	-	-	-	-
		90	14.0±0.0	-	-	-	-	15.0±0.0	9.0±0.0
		120	13.0±0.0	-	-	-	-	18.0±0.0	10.6±1.1
		150	17.5±0.1	-	-	-	-	19.0±0.0	12.0±0.0
<i>C. millenii</i>	2.8	10	-	-	-	-	-	-	-
		30	-	-	-	-	-	-	-
		60	-	-	-	-	-	-	-
		90	-	-	-	-	8.0±0.0	-	8.6±1.1
		120	-	-	-	-	12.0±1.0	11.5±0.0	11.3±0.03
		150	-	-	-	-	18.0±0.0	15.0±0.0	13.6±1.1
<i>C. gigantea</i>	2.5	10	-	-	7.3±0.0	-	-	-	-
		30	-	-	8.0±0.0	-	-	-	-
		60	13.0±0.0	-	-	-	-	-	11.0±0.0
		90	12.0±0.0	-	7.0±0.0	-	-	11.0±0.0	10.3±0.03
		120	14.6±1.1	-	8.0±0.0	-	-	13.0±0.0	13.6±1.1
		150	17.3±0.0	-	8.0±0.0	-	-	13.0±0.0	16.0±0.0
Erythromycin		2.0	20	-	-	-	15	ND	ND
Ketoconazole		2.0	ND	ND	ND	ND	ND	20	25

-: absence of antimicrobial activity

ND: Antimicrobial activity not detected

their habitat. Therefore, the presence of anthraquinones in *C. gigantea* only could be due to environmental conditions. The absence of cardenolides in *Cola millenii* alone is an unusual occurrence; more research is needed to make a meaningful taxonomic deduction of this condition.

Antimicrobial activities shown by the four *Cola* species are in line with previous antimicrobial works on the species of *Cola* (Reid et al., 2005; Adeniyi et al., 2004; Ebana et al., 1991) where *Cola* extracts were found to exhibit important inhibitory activities against the growth of certain bacteria and fungi. The crude ethanolic extract of *C. acuminata*, *C. nitida* and *C. gigantea* showed important activity against *Staphylococcus albus*. The diameters of the zones of inhibition of these extracts were found to be remarkably close to that of the control drug: erythromycin. The MICs were 10mgml⁻¹, 10mgml⁻¹ and 60mgml⁻¹ respectively. However, the leaf ethanol extract of *C. millenii* was inactive against this organism. *C. acuminata* showed the most important activity against *Staphylococcus albus*, *Klebsiella pneumoniae*, *Aspergillus niger* and *Candida albicans*. This is probably due to the strong presence of alkaloids in *C. acuminata* as reported by Adegoke et al., (1968). *C. gigantea* also had a high antimicrobial activity against *Staphylococcus albus*, *Bacillus subtilis*, and on *Aspergillus niger* and *Candida albicans* whereas, *C. nitida* and *C. millenii* had weak inhibitory effects on the growth of all the microorganisms. There is a need for further study to ascertain if the yield in these species would be increased by using stronger fractionating solvents such as ethyl acetone or methyl acetone. These solvents have been reported to be more vigorous than other solvents used in crude extraction of plants (Ajayioba and Fadare, 2006).

An important occurrence is that none of the extracts was effective against *Staphylococcus aureus*. This is in contrast to the observations in some other studies where the tested plants were more active against Gram-positive bacteria (Aladesanmi et al., 2007; Ajaiyeoba and Fadare, 2006; Isu, 2005; Onocha et al., 2003). Generally, the antifungal activities of the extracts as reported in the results were stronger and more pronounced than the antibacterial activities. *C. acuminata* and *C. nitida* showed high inhibitory activity against the two fungi *Aspergillus niger* and *Candida albicans*. The presence of different secondary metabolites in the species of *Cola* is probably offering the therapeutic basis for the antimicrobial activities observed in these species which is in

Table 4: Minimum inhibitory concentration (MIC) of ethanol extracts of four *Cola* species on selected microorganisms

Extract (mgml ⁻¹)	MIC						
	<i>S. albus</i>	<i>S. aureus</i>	<i>B. subtilis</i>	<i>K. pneumoniae</i>	<i>P. aeruginosa</i>	<i>C. albicans</i>	<i>A. niger</i>
<i>C. acuminata</i>							
120	-	+	+	-	+	-	-
90	-	+	+	-	+	+	-
60	-	+	+	+	+	+	+
10	-	+	+	+	+	+	+
<i>C. nitida</i>							
120	-	+	+	+	-	-	-
90	-	+	+	+	-	-	-
60	-	+	+	+	+	+	+
10	-	+	+	+	+	+	+
<i>C. millenii</i>							
120	+	+	+	+	+	-	-
90	+	+	+	+	+	+	-
60	+	+	+	+	+	+	+
10	+	+	+	+	+	+	+
<i>C. gigantean</i>							
120	-	+	-	+	+	-	-
90	-	+	-	+	+	+	-
60	-	+	-	+	+	+	-
10	+	+	-	+	+	+	+

-: No growth observed

+: Growth observed

agreement with other work linking antimicrobial activities with the presence of secondary metabolites (Kisangau, et al., 2007; Kubmamwa et al., 2007; Ajaiyeoba and Sama, 2006; Reid et al., 2005). This claim is further strengthened by the work of Tiew et al. (2003) where the antifungal properties of other members of the Sterculiaceae family were reported. These plants could be a source of new antibiotic compounds. Further work is needed to isolate the secondary metabolites from the extracts studied in order to test specific antimicrobial activity.

In addition, the minimum inhibitory concentration (MIC) of the plants yielded promising results that are worthy of note. *C. acuminata*, *C. nitida* and *C. gigantea* had low MICs of 10mgml⁻¹, 10mgml⁻¹ and 60mgml⁻¹ respectively for *Staphylococcus albus*. This suggests that they can be gainfully employed in the production of antibiotics, as low MICs mean that only a small quantity of the extract will be required to impair bacterial growth. The average minimum MIC of the plants on *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Bacillus subtilis*, *Aspergillus niger* and *Candida albicans* was 90mgml⁻¹, a value which is still low enough to be of great antimicrobial advantage. The closeness observed in the antimicrobial activities demonstrated by *C. nitida* and *C. millenii* as revealed by values obtained for the MIC could also indicate a close relationship between the two species.

In conclusion, the occurrence of a similar pattern of secondary metabolites in three out of the four species of *Cola* studied is suggestive of an important trend in the species. Further studies on structural elucidation of the compounds may offer some information which may become very useful in the knowledge of the natural relationship of the four plants. The ethanol crude extract of *C. acuminata* had the most important effect against both bacteria and fungi. Ethanol crude extracts of the other three species also had important effects on some of the microorganisms. Therefore, the plants are justified in their ethnomedicinal uses in the treatment of certain diseases, especially fungal diseases. The comparison of data obtained suggests a close relationship between *C. gigantea* and *C. acuminata* and also between *C. nitida* and *C. millenii*. Based on the presence of the

typical secondary metabolites in the ethanol leaf extracts of the four species together with the antimicrobial activities demonstrated against various organisms, our study has highlighted the possible usefulness of phytochemical and antimicrobial studies as taxonomic tools in evaluating closeness among four species of the genus *Cola*. This provides evidence for further research in the chemical profile of the genus. It is important to mention that *C. acuminata* and *C. gigantea* gave the best all-round results.

Acknowledgements

We are grateful to Mr T. K. Odewo of the Forest Herbarium, Ibadan for helping with the identification and authentication of the plants and also to the anonymous reviewer for helpful comments on the earlier version of the manuscript.

References

1. Adegoke, E. Akinsanya, A., Nagu, A. (1968). Studies of Nigerian Medicinal Plants: A preliminary survey of Plant Alkaloids. *J. West Afri. Sci. Assoc.* **13**: 13-39.
2. Adeniyi, B. A., Groves, M. J., Gangadharam, P. R. J. (2004). In vitro anti-mycobacterial activities of three species of Cola plant extracts (Sterculiaceae). *Phytother. Res.* **18** (5): 414-418.
3. Ajaiyeoba, E. O and Fadare, D. A. (2006). Antimicrobial potential of extracts and fractions of the African Walnut – *Tetracarpidium conophorum*. *Afr. J. Biotechnol.* **5** (22): 2322-2325.
4. Ajaiyeoba, E. O and Sama, W. (2006). Phytochemical and antimicrobial studies of *Capparis thoningii* and *Capparis tomentosa*. *Pharmacognosy Magazine*, **2**(6): 119 –122.
5. Ajaiyeoba, E.O., P. A. Onocha, S. O. Nwozo and W. Sama (2003). Antimicrobial and cytotoxicity evaluation of *Buchholzis coricea* stem bark. *Fitoterapia*, **70**: 184-186.
6. Aladesanmi, A. J., Iwalewa, E. O., Adebajo, A. C., Akinkunmi, E. O., Taiwo, B. J., Olorunmola, F. O. and Lamikanra, A. (2007). Antimicrobial and Antioxidant activities of some Nigerian medicinal plants. *Afri. J. Trad. CAM* **4** (2): 173-184
7. Benjamin, L. T., Rogers, A. M, Rosenbaum, A. (1991). “Coca-cola”, Caffeine, and Mental Deficiency - Harry Hollingworth and the Chattanooga trial of 1991”. *J. Histol. Behavioural Sci.* **27** (1): 42-45.
8. Blades, M. (2000). Functional foods or nutraceuticals. *Nutrition and Food Sci.* 30 (2): 73-75
9. Bradley, P. R. (ed.). (1992). *British Herbal Compendium*. Vol. 1. Bournemouth: British Herbal Medicine Association.
10. Braga, M. R., Adiar, P. M., Marabesi, M. A. And de-Godoy, J. R. L. (2006). Effects of elevated CO₂ on the phytoalexin production of two soybean cultivars differing in the resistance to stem canker disease. *Environ. Exp. Bot.* **58** (1-3): 85-92.
11. Burkill, H. M. (1995). *The Useful Plants of West Tropical Africa*, 2nd Edition, Vol 3:11. Royal Kew Botanical Gardens, Kew. London. pp. 522 – 527.
12. Chevalier, A. and Perrot, A. (1911). “Les vegetaux utiles de l’afrique tropicale française”. *Cashier d’Etudes Africanes*, **29**:101-107
13. Cybulskill, W. J., Peterjohn, W. T. and Sullivan, J. H. (2000). The influence of ultraviolet-B radiation on tissue quality and composition of loblolly pine (*Pinus taeda* L.) needles. *Environ. Exp. Bot.* **44** (3): 231-241
14. Daels-Rakotoarison, D. A., Kouakoua, G., Gressiera, B., Dinea, T., Bruneta, C., Luyckxa, M., Bailleul, F. and Trotin, F. (2003). Effects of a caffeine-free *Cola nitida* nuts extract on elastase/alpha-1-proteinase inhibitor balance. *J. Ethnopharmacol.* **89** (1): 143-150
15. Duke, J. A. (2001). *Handbook of Nuts*. CRC Press, Boca Raton, FL. pp. 343.
16. Ebana. R. U. B, Ekpe B. E, Madunagu, B. E. and Otung, I. N. (1991). Microbiological Exploitation of Cardiac Glycosides and Alkaloids from *Garcinia kola*, *Borreria ocymoides*, *Kola nitida* and *Citrus aurantifolia*. *J. Appl. Bacteriol.* **71**: 398-401.
17. Folkers, A., Hüve, K., Ammann, C., Dindorf, T., Kesselmeier, J., Kleist, E., Kuhn, U., Uerlings, R. And Wildt, J. (2008). Methanol emissions from deciduous tree species: dependence on temperature and light intensity. *Plant Biol.* **10** (1): 65-75
18. GRIN, (2007). *Germplasm esources Information Network*. Beltsville, Maryland
19. Irvine, F. R. (1961). *Wood Plants of the Ghana*. Oxford University Press; **17**: 498-502.
20. Isu, N. R. (2005). Antibacterial effect of *Aframomum melegueta*, *Xylopia aethiopicum* and *Ocimum viride*. *Nig. J. Nat. Prod. Med.* **9**: 22-25
21. Jayeola, O. C. (2001). Preliminary studies on the use of kolanuts (*Cola nitida*) for soft drink production. *J. Food Technol. Afr.* **6** (1): 25-26

22. Keay, R. W. J. (1989). *Trees of Nigeria*. Clarendon Press. Oxford. pp. 131-135.
23. Kim, K. (2001). *Encyclopedia of Alternative Medicine*. **22**: 203-204.
24. Kisangau, D. P., Hosea, K. M., Joseph, C. C. and Lyaruu, H. V. M. (2007). In vitro antimicrobial assay of plants used in Traditional Medicine in Bukoba Rural District, Tanzania. *Afri. J. Trad. CAM*. **4 (4)**: 510 – 523
25. Kubmamwa, D, Ajoju, G. A, Enwerem, N. N. and Okarie, D. A. (2007). Preliminary phytochemical and Antimicrobial Screening of 50 Medicinal Plants from Nigeria. *Afr. J. Biotechnol.* **6 (14)**: 1690-1696.
26. Kuoame, C. and Sacande, M. (2006). *Cola nitida* (Vent.) Schott & Endl. Seed Leaflet 111.
27. Newall, C. A., Anderson, L. A., Phillipson, J. D. (1996). *Herbal Medicines: A guide for Health Professionals*. London: The Pharmaceutical Press. pp. 84.
28. Odugbemi, T. (2006). *Outlines and Pictures of Medicinal plants from Nigeria*. University of Lagos Press; 10. pp. 158.
29. Oliver-Bever, B. (1986). *Medicinal plants in Tropical West Africa*, Cambridge University Press, Great Britain. pp.375.
30. Oliver, B. (1960). *Medicinal Plant in Nigeria*. Nigerian College of Arts, Science and Technology. Ibadan.
31. Olorode, O. (1984). *Taxonomy of West African Flowering Plants*. Longman Group, United Kingdom. pp. 180
32. Onocha, P. A., Opegbemi, A. O., Kadri, A. O., Ajayi, K. M. And Okorie, D. A. (2003). Antimicrobial evaluation of Nigerian Euphorbiaceae plants 1. *Phyllanthus amarus* and *Phyllanthus muellerianus* leaf extracts. *Nig. J. Nat. Prod. Med.* **7**: 9-12
33. Opeke, L. K. (1992). *Tropical Tree Crops*. Spectrum books Ltd, Ibadan. pp. 180-200.
34. Purseglove, J. W. (1968). *Tropical Crops-Dicotyledon*. Longman, Green and Co. Ltd, London, **8**: 62-63
35. Ratsch, C. (2005). *The Encyclopedia of Psychoactive plants: Ethnopharmacology and its applications*. Foreword Publisher.
36. Reid, K. A., Jager, A. K., Light, M. E., Mulholland, D. A. and Staden, J. V. (2005). Phytochemical and Pharmacological Screening of Sterculiaceae species and Isolation of antibacterial Compounds. *J. Ethnopharmacol.* **97 (2)**: 285-291.
37. Russell, T. A. (1955). The kola of Nigeria and the Cameroons. *Trop. Agr., Trinidad*, **32**: 210-240.
38. Sahm DF, JA Washington (1990). Antibacterial susceptibility Test Dilutions Methods: In: *Manuals of Clinical Microbiology*. Lennette E.H 5th Edition, Am. Soc. Microbiol. Washington DC, pp. 1105-1116.
39. Shen, H., Tang, Y., Muraoka, H. and Washitani, I. (2008). Characteristics of leaf photosynthesis and simulated individual carbon budget in *Primula mutans* under contrasting light and temperature conditions. *J. Plant Res.* **121**: 191-200
40. Steinegger, E. and Hansel, R. (1992). *Pharmakognosie*. 5th ed. Berlin-Heidelberg. Springer Verlag.
41. Stich, B. (1932) Subacute bacterial endocarditis due to *Staphylococcus albus*. *Arch. Intern. Med.* **49 (4)**: 666-670
41. Tiew, P., Ioset, J. R., Kokpol, U., Chavasiri, W., Hostettman, K. (2003). Antifungal, Antitoxidant And Larvicidal Activities of Compounds Isolated From The Heartwood of *Mansonia gagei*. *Phytother. Res.* **17 (2)**: 190-193.
42. Trease, G. E. and Evans, W. C. (1989). *Trease & Evans' Pharmacognosy* 13th Edition. Bailliere Tindale London. pp. 832.
43. Tsukaya, H., Tsujino, R., Ikeuchi, M., Isshiki, Y., Kono, M., Takeuchi, T. and Araki, T. (2007). Morphological variation in leaf shape in *Ainsliaea apiculata* with special reference to the endemic characters of populations on Yakushima Island, Japan. *J. Plant Sci. Res.* **120 (3)**: 351-35
44. Wilson, T. S and Stuart, R. D. (1965). *Staphylococcus albus* in wound infection and septicemia. *Canad. Med. Ass. J.* **93(1)**: 8-16.