

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/283034736>

Flavonoids from the roots of *Dioclea reflexa* (Hook F.)

Article in *Bulletin of the Chemical Society of Ethiopia* · October 2015

DOI: 10.4314/bcse.v29i3.12

CITATIONS

4

READS

414

4 authors, including:



Abdulkabir Oladimeji

University of Ibadan

17 PUBLICATIONS 44 CITATIONS

[SEE PROFILE](#)



Ibrahim Adebayo Oladosu

University of Ibadan

63 PUBLICATIONS 195 CITATIONS

[SEE PROFILE](#)



Zeeshan Ahmed

University of Karachi

20 PUBLICATIONS 102 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



bio-prospecting for anti-tuberculosis lead compounds from Nigerian medicinal plants [View project](#)



Anti-mycobacterium tuberculosis from insect fungus. [View project](#)

FLAVONOIDS FROM THE ROOTS OF *DIOCLEA REFLEXA* (HOOK F.)

Abdulkabir O. Oladimeji^{1,2}, Ibrahim A. Oladosu^{1*}, Mohammad S. Ali³ and Zeeshan Ahmed³

¹Organic and Medicinal Unit, Department of Chemistry, University of Ibadan, Nigeria

²Department of Chemical and Geological Sciences, Al-Hikmah University, Ilorin, Nigeria

³International Centre for Chemical and Biological Sciences (ICCBS), Karachi, Pakistan


(Received May 1, 2014; revised June 7, 2015)

ABSTRACT. Reflevone (**1**), a new flavone, together with mearnsetin (**2**), 7,4'-dihydroxyflavone (**3**) and two known phytosterols, β -sitosterol (**4**) and stigmasterol (**5**) were isolated from ethanol soluble part of the roots of *Dioclea reflexa* (Hook F.). Their structures were elucidated by analysis of spectroscopic data involving 1D and 2D NMR, MS, IR and UV experiments. The crude roots extract showed significant activity with $IC_{50} = 5.18 \mu\text{g/mL}$ against the prostate cancer cell line.

KEY WORDS: Phytochemical analysis, *Dioclea reflexa*, Fabaceae, Cytotoxic activity, Flavonoid

INTRODUCTION

Dioclea (Leguminosae-Fabacea) is a small genus containing 40 species, widely distributed in Africa and South America [1]. The pharmacological activities of this genus include anticonvulsant, anxiolytic [2], antioxidant [3], antiarrhythmogenic [4], immunodulatory [5], analgesic and central antinociceptive effects [6], which have been identified with *Dioclea* plants. *Dioclea reflexa* (Hook F.) is a woody climber, which produces flowers and pods on long rope-like stems that hang from the forest canopy. It is known as 'Agbaarin' in Western Nigeria. The stem extract of this species is used by the local populace as an anti-tuberculosis (TB), and anti-inflammatory agents; and also in the treatment of rheumatism and itching [7, 8]. Previous studies of the seed have led to the isolation of dioclimidazole and two steroids: taraxasterol and stigmasterol [7, 9]. In our earlier work on the root of *D. reflexa*, we characterized 15 volatile compounds comprising monoterpenes and diterpenes [8]. Although, several secondary metabolites have been isolated from the seeds [7, 8] of *D. reflexa*, none have been reported from the roots of this species. In our search for the active constituents of *D. reflexa*, we hereby report an investigation of the cytotoxic activity of the crude extracts of leaves, stems and roots of *D. reflexa* against prostate cancer cell (PC-3). This led to the isolation and characterization of a new flavone, reflevone (**1**), together with mearnsetin (**2**), 7,4'-dihydroxyflavone (**3**), β -sitosterol (**4**) and stigmasterol (**5**) from roots extract. This is the first reported isolation of compounds **1-3** from this species.



*Corresponding author. E-mail: dosufau@gmail.com

Compound	R ₁	R ₂	R ₃	R ₄	R ₅
1	H	H	OH	CH ₃	OH
2	OH	OH	OH	CH ₃	OH
3	H	H	H	H	H

EXPERIMENTAL

General experimental procedures

The ¹H-NMR spectra (δ ppm, J in Hz) were recorded in deuteriated acetone or methanol on a Bruker AM-400 spectrometer (400 MHz), while ¹³C-NMR spectra were recorded on the same instrument using the same solvent at 100 MHz. IR and UV were recorded using a JASCO 302-A and SHIMADZU UV-240 spectrometers respectively. HR-EIMS were carried out using MAT 95XP spectrometer. For the detection of samples on TLC plates, ceric sulfate spray reagent and UV light were used. TLC was carried out to check the purity of samples on pre-coated Merck silica gel F₂₅₄ aluminium sheets (0.25 mm thickness). Column chromatography was performed on silica gel (70-230 mesh) and Sephadex LH-20 (Fluka 25-100 mm). Melting points were determined on a Buchi 510-K melting point apparatus and were uncorrected. Enzyme-linked immunosorbent assays were recorded on Spectra Max plus 384, Molecular Devices, CA, USA, reader.

Plant material

The whole plant of *Dioclea reflexa* was collected from a farm land at Eruwa in Ibarapa East Local Government area of Oyo state, Nigeria, in May, 2011, and authenticated by Mr Bolu Ajayi of Herbarium Section, Plant Biology Department, University of Ilorin, Nigeria and a voucher specimen (UIL-0027) was deposited at the herbarium of the University.

Extraction and isolation

The air-dried and powdered roots of *D. reflexa* (1 kg) were extracted with 5 L of ethanol at room temperature for two weeks. Evaporation of the solvent under vacuo yielded crude root ethanol extract (85 g). The same extraction procedure was adopted for the leaves and stem ethanol extracts. These extracts were first subjected to preliminary anticancer assay. The crude root ethanol extract was then subjected to silica gel column chromatography, eluted with gradient of n-hexane, dichloromethane, ethyl acetate and methanol in increasing polarity. The 332 fractions (100 mL/vial) collected were combined according to TLC profile.

The combined fractions 205-241 (1.2 g), eluted with n-hexane/EtOAc (R_f 55:45) was further purified by column chromatography using n-hexane and EtOAc gradient to give **1** (7 mg), in addition to four semi-pure sub-fractions labeled F₁-F₄.

Sub-fraction F₁ (262 mg) was subjected to repeated column chromatography in Sephadex LH-20 using 100% MeOH elution, which led to the isolation of compound **2** (5 mg) as yellow powder. Similarly, fraction labelled F₂ yielded compound **3** (5 mg). Compounds **4** (9.0 mg) and **5** (4.8 mg) were also obtained from repeated column chromatography over silica gel and Sephadex of sub-fractions F₃ and F₄ respectively.

Physical and spectral data of the compound (1)

7-Hydroxy-2-(3,5-dihydroxy-4-methoxyphenyl)-4H-chromen-4-one, C₁₆H₁₂O₆; yellow crystals, mp 220 °C. UV λ_{max} nm (log ε) = 212 (1.45), 311 (0.9); IR (KBr): 3377 (OH), 1676 (C=O), 1570 (C=C) cm⁻¹; EIMS *m/z* (relative intensity %): 53 (48), 77 (22), 108 (20), 121 (40), 137 (40), 149 (29), 164 (8), 200 (8), 229 (100), 257 (96), 300 (89). HREIMS = 300.0621 (calculated = 300.2700 for C₁₆H₁₂O₆). ¹H-NMR (CD₃COCD₃, 400 MHz) (Table 2). ¹³C-NMR (CD₃COCD₃, 100 MHz) (Table 2).

Physical and spectral data of the compound (2)

3,5,7-Trihydroxy-2-(3,5-dihydroxy-4-methoxyphenyl)-4H-chromen-4-one, C₁₆H₁₂O₈; yellow powder, mp 208-210 °C. UV λ_{max} nm (log ε) = 212 (1.268), 262 (0.749), 365 (0.845); IR (KBr): 1676 (C=O ketone), 1659 (ring C=C), 1597, 1459, 1452, 849 cm⁻¹; EIMS *m/z* (relative intensity %): 332.1 (91), 317 (M⁺, 100), 289 (20), 261 (48), 233 (11), 205 (10), 177 (6), 153 (20), 108 (12), 77 (32). HREIMS = 332.0574 (calcd. 332.25287 for C₁₆H₁₂O₈). ¹H-NMR (CD₃COCD₃, 400 MHz) (Table 3). ¹³C-NMR (CD₃COCD₃, 100 MHz) (Table 3).

Physical and spectral data of the compound (3)

7-Hydroxy-2-(4-hydroxyphenyl)-4H-chromen-4-one, C₁₅H₁₀O₄; yellow powder, mp 315 °C. UV λ_{max} nm (log ε) = 213 (1.627), 230 (1.362), 328 (1.677); IR (KBr) = 3619 (Ar-OH), 1676 (ketones C=O), 1649 (ring C=C) cm⁻¹; EIMS *m/z* (relative intensity %): 254.2 (M⁺, 100), 237 (5.5), 226 (41), 136 (9), 137 (40), 118 (24), 86 (39), HREIMS = 254.0567 (calcd. 254.24425 for C₁₅H₁₀O₄). ¹H-NMR (CD₃OD, 400 MHz) (Table 3). ¹³C-NMR (CD₃OD, 100 MHz) (Table 3).

In vitro cytotoxicity assay

The cytotoxic activity against Prostrate Cancer (PC-3) cells using test extract was performed in 96-well flat-bottomed micro plates by employing 3-[4,5-dimethylthiazole-2-yl]-2,5-diphenyl-tetrazolium bromide (MTT) colorimetric assay [10]. The PC-3 cells were grown in Dulbecco's Modified Eagle Medium, supplemented with 5% fetal bovine serum (FBS) and antibiotics (100 µg/mL of penicillin and 100 µg/mL of streptomycin) under a 5% CO₂ humidified atmosphere at 37 °C incubation. Cells culture with a density of 1 x 10⁵ cells/mL were exposed to plant extract dissolved in DMSO at various concentrations of 1-30 µM. After 48 h, cells were washed 3 times with FBS, and then 200 µL of 0.5 mg/mL MTT solution in FBS was added and incubated for 4 h. Finally, 100 µL of DMSO was added to each well. In the experiment, Doxorubicin was used as positive control. The optical density was measured spectrophotometrically at 570 nm in an enzyme-linked immunosorbent assay reader (Spectra Max plus 384, Molecular Devices, CA, USA). The 50% inhibition concentration (IC₅₀) was calculated as the compound concentration required to reduce MTT signals by 50% compared with untreated control cultures.

RESULTS AND DISCUSSION

The IC₅₀ values of various parts extract of *D. reflexa* when tested against PC-3 cell lines are presented in Table 1. The roots extract showed cytotoxic activity against the cell lines (IC₅₀ 5.18 µg/mL) when compared to the positive control, doxorubicin (2.8 µg/mL). This strong cytotoxic activity prompted further investigation of the roots ethanol extract of *D. reflexa*. The roots extract was then subjected to repeated column chromatography which yielded compound **1**.

Table 1. Results of anticancer assay of various parts of *D. reflexa* extracts.

Extracts	IC ₅₀ ± SEM (µg/mL) PC3
Leaves	NA
Stems	11.42 ± 0.7
Roots	5.18 ± 0.3
Doxorubicin	2.8 ± 0.12

PC3, human prostate cancer cell; NA, not active; SEM, standard error of mean.

Compound **1** was isolated as a yellow powder. The molecular formula was established as C₁₆H₁₂O₆ (300.0621) corresponding to 11 units of unsaturation based on the HREIMS, ¹³C and DEPT NMR data (Table 2). The UV spectrum of **1** showed maxima at λ_{max} (logε) nm 212 (1.45)

and 311(0.9) similar to those of flavones skeleton with two absorption bands. The IR spectrum suggested the presence of hydroxyl (3377 cm^{-1}), conjugated carbonyl (1776 cm^{-1}) and aromatic system (1570 cm^{-1}).

Table 2. ^1H and ^{13}C NMR spectra data of compound **1** (δ values; 400 MHz and 100 MHz, acetone- d_6).

Position	^1H δ (integration, multiplicity, J in Hz)	^{13}C	DEPT	HMBC
2	-	163.30	C	
3	6.54 (1H, s)	107.03	CH	C-2, C-4, C-10, C-1'
4	-	177.15	C	
5	7.95 (1H, d, $J = 8.8$ Hz)	127.66	CH	C-4, C-7, C-8, C-9
6	6.97 (1H, dd, $J = 2.4, 8.8$ Hz)	115.43	CH	C-8, C-10
7	9.64 (1H, br s, OH)	163.30	C	
8	7.02 (1H, d, $J = 2.4$ Hz)	103.47	CH	C-6, C-7, C-9, C-10
9	-	158.74	C	
10	-	117.95	C	
1'	-	128.26	C	
2'/6'	7.07 (1H, s)	106.58	CH	C-2, C-1', C-3', C-4'
3'/5'	8.48 (s, OH)	151.83	C	
4'	-	139.00	C	
OCH ₃	3.88 (3H, s)	60.74	CH ₃	C-4'

The analysis of proton NMR spectrum of **1** revealed the presence of one methoxy [δ_{H} 3.88 (s); δ_{C} 60.7], one *meta*-coupled [δ_{H} 7.02, 1H, d, $J = 2.4$ Hz], one *ortho*-coupled [δ_{H} 7.95, 1H, d, $J = 8.8$ Hz], two supposed *meta*-coupled but appeared as singlet each because they are symmetrical [δ_{H} 7.07, 2 x 1H, s] and one double-doublet *ortho-meta*-coupled aromatic protons [δ_{H} 6.97, 1H, dd, $J = 2.4$ and 8.8 Hz], three phenolic hydroxyl groups of which two are symmetrical [δ_{H} 8.48, 8.48, 9.64 (1H each, br s)], and an olefinic hydrogen at δ_{H} 6.54, 1H, s.

The ^{13}C -NMR (Table 2), DEPT and HMBC spectra showed resonances for all 16 carbons comprising one methoxy, six methine and nine quaternary carbons. The most downfield shifted peak at δ_{C} 177.15 was assigned to carbonyl group (C-4). Two peaks at δ_{C} 106.58 and δ_{C} 151.83 showed double intensities and were assigned to C-2'/6' and C-3'/5', respectively [11, 12]. The HMBC spectrum of **1** showed cross-peaks from H-2' to C-3', C-4', and C-2; from H-3 to C-4, C-2 and C-1' confirming the assignments for C-2, H2'/6' and the B-ring. In addition, the methoxy at δ_{H} 3.88 showed correlation with carbon resonance at δ_{C} 139.0 on the HMBC spectrum and was assigned C-4' (Figure 1).

Furthermore, the ^1H - ^1H COSY spectrum demonstrated correlations of H-5 to H-6 (Figure 1). The spectral data were in accordance with reported data on flavonoids of 7-hydroxy A-ring [11, 12].





Figure 1. HMBC and COSY connectivities of **1**. Key COSY (→) and HMBC (→).

The 2H singlet at δ 7.07, 2H broad singlet at δ 8.48 and 3H singlet of methoxy at δ 3.88 in ring B were assigned to H-2'/6', OH-3'/5' and H-4', respectively; this suggested a 3',4',5'-trioxygenated symmetrical substituted B-ring, typical of mearnsetin B-ring [13]. In HMBC, C-4 showed long-range connectivity to the ^1H peak at 7.95 ppm which was directly attached to the ^{13}C peak at 127.66 ppm (HSQC spectrum) and was assigned as C-5. In addition, the ^1H at 7.95 ppm showed long-range coupling to ^{13}C signals at 163.3 ppm and 158.74 ppm, these were assigned as C-7 and C-9, respectively. The pronounced upfield shift of ^{13}C peak at 163.3 ppm further supported the presence of a hydroxyl at C-7. The COSY spectrum as well as the HMQC spectrum confirmed the proposed structure (Figure 1). The complete assignments of the ^1H and ^{13}C -NMR data are listed in Table 2. These observations suggested that compound **1** is a 7,3',5'-trihydroxy-4'-methoxyflavone. Extensive literature search further revealed that compound **1** is new and has not been reported in the chemical literature, hence it was given the trivial name *reflevone*.



Scheme 1. Fragmentation pattern for compound **1**.

The retro-Diels-Alder fragments of ring C led to diagnostic peaks at m/z 137 and 164, supporting the presence of one hydroxyl in ring A, a methoxy and two hydroxyls in ring B

(Scheme 1). Another diagnostic fragment at m/z 149 ($C_8H_5O_3$) indicated methoxy moiety at C-4' which was demethylated (Scheme 1). Other significant peaks in addition to molecular ion at m/z 300.1 are 285 [M^+-CH_3], 257, 229 and 121.

Table 3. 1H (400 MHz) and ^{13}C (100 MHz) NMR Spectra data of **2** (in CD_3COCD_3) and **3** (in CD_3OD).

Position	1H δ (integration, multiplicity, J in Hz)	^{13}C	DEPT	1H δ (integration, multiplicity, J in Hz)	^{13}C	DEPT
2	-	146.16	C	-	166.02	C
3	-	137.52	C	-	105.13	CH
4	-	176.73	C	-	180.30	C
5	12.10 (1H, s, OH)	162.3	C	7.96 (1H, d, J = 8.8)	127.75	CH
6	6.26 (1H, d, J = 2 Hz)	99.18	CH	6.91 (1H, dd, J = 2, 8.8)	116.32	CH
7	-	165.14	C	-	165.00	C
8	6.52 (1H, d, J = 2 Hz)	94.44	CH	6.97 (1H, d, J = 2)	103.50	CH
9	-	157.79	C	-	159.50	C
10	-	104.17	C	-	114.30	C
1'	-	127.31	C	-	123.50	C
2'/6'	7.38 (2H, s)	108.31	CH	7.86 (2H, dt, J = 2.8, 9.2)	129.39	CH
3'/5'	8.36 (2H, s, OH)	151.31	C	6.91 (2H, m, J = 2.4, 8.8)	117.01	CH
4'	-	138.02	C	-	162.50	C
OCH ₃	3.88 (3H, s)	60.69	CH ₃	-	-	-




Figure 2. HMBC and COSY connectivities of **2**. Key COSY (—) and HMBC (→).

Compound **2**, obtained as a yellow powder was assigned the molecular formula $C_{16}H_{12}O_8$ with molecular ion peak at m/z 332 from HR-EIMS and supported by its NMR spectra data (Table 3). The UV spectrum in methanol showed maximum absorptions at 365, 260 and 212 nm. The 1H -NMR spectrum (Figure 2) showed two meta coupled 1H doublets at 6.26 ($J = 2$ Hz) and 6.52 ($J = 2$ Hz) assigned to H-6 and H-8 respectively corresponding to δ_C 99.18 ppm and δ_C 94.44 ppm respectively in the HSQC spectrum, characterised the H-6 and H-8 of a flavonoid 5,7-dihydroxyl A-ring [14]. The ^{13}C -NMR spectrum revealed all the sixteen carbons which were deduced as one methoxyl, four methines and eleven quaternary carbons. The broad singlet 1H peaks at δ_H 8.36 ppm which was not directly attached to any carbon in HSQC, was assigned to 3'/5'-OH. In the 1H -NMR spectrum, 3H singlet at δ_H 3.88 revealing a methoxyl group correlated with the carbon resonance at 138.02 (C-4') on the HMBC spectrum. The 2H singlets at δ_H 7.38 was attributed to the one proton each at 2' and 6' positions of 3',4',5' tri-O-substituted B ring. Long-range COSY spectrum in conjunction with the HMBC spectrum allowed the assignments of H-6 and H-8 (Figure 2). A singlet at δ_H 12.1 corresponding to ^{13}C peak at δ_C 162.3 ppm in HSQC, implied the presence of an OH at C-5, since H-bonding with the C=O (C-4) causes this

type of substituted flavonoid to give signal between 159.8 and 162.3. Also, the signals for C-5 unsubstituted flavonoids are usually found between 106.9 and 127.75 [12]. Thus compound **2** was identified as 3,5,7,3',5'-pentahydroxy-4-methoxyflavone (mearnsetin) by comparison of its spectra data with the literature [13, 14].





Figure 3. HMBC connectivities of **3**. Keys: COSY (—) and HMBC (---).



Scheme 2. Fragmentation pattern of compound **3**.

The $^1\text{H-NMR}$ spectrum of compound **3** displayed two 1H doublets at δ 6.97 and δ 7.96 assigned to H-8 and H-5 respectively. A 3H multiplet at δ 6.91 was assigned to H-6 and two equivalent aromatic protons at H-3'/5'. The $^{13}\text{C NMR}$, DEPT and HMBC experiments showed signals for 15 carbons which were deduced as: seven quaternary and eight methine carbons. The

various connectivities in the molecule were determined through HMBC and COSY experiments, which concluded the position of OH at C-7 and C-4' (Figure 3). The molecular formula of compound **3** (C₁₅H₁₀O₄) was deduced from the high resolution electron impact mass spectrum (HR-EIMS). The Retro-Diels-Alder (RDA) fragmentation of the heterocyclic ring produced ions at m/z 137[A+1]⁺ and m/z 118 [B]⁺ (Scheme 2) which indicated that A-ring and B-ring possessed OH substituent each. Other significant peaks in addition to the molecular ion at m/z 254 are 237 [M-OH]⁺, 226 [C₁₄H₁₀O₃], 137 [C₇H₅O₃] and 108 [C₆H₄O₂]. The comparison of the corresponding UV, EIMS, ¹H and ¹³C NMR data (Table 3) with those reported in literature [11, 15-17] led to its identification as 7,4'-dihydroxyflavone. Compounds **4** and **5** were identified as β-sitosterol and stigmaterol respectively [18]. This is the first reported isolation of compounds **1-3** from this species.

In conclusion, the present study has shown that roots extracts of *D. reflexa* could be a lead in discovery of anti-cancer drug as demonstrated by its cytotoxic effect. Purification of the roots extracts led to the isolation of a new flavone compound and four others. Further studies are suggested on screening the isolated compounds against PC-3 cell lines, particularly the new compound, refflavone.

ACKNOWLEDGMENTS

The authors are grateful to Third World Academy of Science (TWAS) and International Centre for Chemical and Biological sciences (ICCBS) for fellowship support to A.O. Oladimeji. We are indebted to Mr. Bolu Ajayi of Herbarium Section, Plant Biology Department, University of Ilorin for his assistance in the identification and authentication of the plant.

REFERENCES

- Burkill, H.M. *The Useful Plants of West Tropical Africa*, 2nd ed., Royal Botanic Gardens: UK; **1984**;
- De Almeida, E.R.; Xavier, H.S.; Chaves, T.M.; Couto, G.B.L.; Aragao-Neto, A.C.; Silva, A.R.; Da Silva, L.L.S. *Int. J. App. Res. Nat. Prod.* **2010**, 2, 44.
- Barreiros, A.L.B.S.; David, J.P.; Queiroz, L.P.; David, J.M. *Phytochemistry* **2000**, 55, 805.
- Vianna, H.R.; Cortes, S.F.; Ferreira, A.J.; Carpettini, L.S.A.; Schnitt, M.; Almeida, A.P.; Massensini, A.R.; Lemos, V.S. *Planta Med.* **2006**, 72, 300.
- Alves, C.Q.; Kijjoa, A.; David, J.M.; David, J.P. *Planta Med.* **2010**, 76-P368. DOI:10.551s-0030-126466.
- Almeida, E.R.; Almeida, R.N.; Navairo, D.S.; Bhattacharya, J.; Silva, V.A.; Birnbaum, J.S.P. *J. Ethnopharmacol.* **2003**, 88, 1.
- Faleye, F.J. *J. Pharm. Sci. Innov.* **2012**, 1, 89.
- Oladosu, I.A.; Oladimeji, A.O. *Elixir Org. Chem.* **2012**, 45, 7738.
- Oladosu, I.A.; Echeme, J.O.; Zubair, M.F. *Fitoterapia* **2006**, 77, 571.
- Mosmann, T. *J. Immunol. Methods* **1983**, 65, 55.
- Park, Y.; Moon, B.; Lee, E.; Lee, Y.; Yoon, Y.; Ahn, J.; Lim, Y. *Magn. Reson. Chem.* **2007**, 45, 674.
- Park, Y.; Lee, Y.; Kim, H.; Lee, Y.; Yoon, Y.; Moon, B.; Chong, Y.; Ahn, J.; Shim, Y.; Lim, Y. *Bull. Korean Chem. Soc.* **2006**, 27, 1537.
- Abbas, F.A.; Al-Massarany, S.M.; Khan, S.; Al-Howiriny, T.A.; Mossa, J.S.; Abdourasheed, E.A. *Nat. Prod. Res.* **2007**, 21, 383.
- Gluchoff-Fiasson, K.; Fenet, B.; Leclerc, J.; Lussignol, M.; Jay, M. *Chem. Pharm. Bull.* **2001**, 49, 768.
- Bickoff, E.M.; Livingston, A.L.; Witt, S.C. *Phytochemistry* **1964**, 9, 523.
- Maxwell, C.A.; Hartwig, U.A.; Joseph, C.M.; Phillips, D.A. *Plant Physiol.* **1989**, 91, 842.
- Meskuriyen, D.; Cordell, G.A. *J. Sci. Soc. Thailand* **1988**, 14, 24.
- Ngnokam, D.; Massiot, G.; Nuzillard, J.M.; Tsamo, E. *Bull. Chem. Soc. Ethiop.* **1994**, 8, 15.