

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/244944245>

Antioxidant and antibacterial activities of Trema orientalis Linn: an indigenous medicinal plant of Indian subcontinent

Article in *Oriental Pharmacy and Experimental Medicine* · December 2008

DOI: 10.3742/OPEM.2008.8.4.395

CITATIONS

9

READS

376

1 author:



Sarder N. Uddin

MSU, USA

77 PUBLICATIONS 510 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Neural circuit development and processing of sensory information [View project](#)



Autism in Bangladesh [View project](#)



Antioxidant and antibacterial activities of *Trema orientalis* Linn: an indigenous medicinal plant of Indian subcontinent

Sarder Nasir Uddin^{1,2,*}

¹Department of Biology and Chemistry, Graduate School of Medicine, Yamaguchi University, Yamaguchi-735-8512, Japan; ²Biotechnology and Genetic Engineering Discipline, Khulna University, Bangladesh

Received for publication October 25, 2007; accepted August 08, 2008

SUMMARY

Methanol and aqueous extract of leaves of *Trema orientalis* Linn. were subjected to the potential antioxidant and antibacterial activities. The pharmacological interest of this plant coupled with traditional use (antidiarrhoeal, antiseptic, analgesic etc) prompted to test for antioxidant and antibacterial activities. The antioxidant potential of the methanolic extract was determined on the basis of their scavenging activity of the stable 1,1-diphenyl-2-picryl hydrazyl free radical. IC₅₀ of the methanol extract of *T. orientalis* was 110.25 µg/ml which indicated the strong antioxidant activity of the plant. However the aqueous extract showed mild antioxidant activity. In case of antibacterial activities test, the extract was subjected for its effectiveness against both Gram-positive and Gram-negative bacteria in agar diffusion method. The zones of inhibition produced by the crude methanol and aqueous extract against few sensitive strains were measured and compared with those of standard antibiotic Gentamycin. It is evident that both extracts are active against the bacteria at low concentrations. The obtained results provide a support for the use of this plant in traditional medicine and suggest its further advance investigation.

Key words: Antioxidant; Antibacterial; *Trema orientalis* Linn.; Activities; Extract

INTRODUCTION

Bangladesh possesses rich floristic wealth and diversified genetic resources of medicinal plants. It has a widely ranging tropical and the agro climatic conditions, which are conducive for introducing and domesticating new and exotic plant varieties. The use of the plants, plant extracts and pure compounds isolated from natural sources provided the foundation to modern pharmaceutical compounds. *Trema orientalis* Linn. is one of the common

medicinal plant grown in Indian subcontinent. Different parts of this plant have been used in traditional medicine.

T. orientalis (Bengali name- Jibon or Chikor) is a tree and belongs to the Ulmaceae family. The plant is distributed in almost all districts of Bangladesh and is used in traditional medicine by the rural people and possesses various interesting pharmacological activities (Uddin *et al.*, 2008). The root of the plant is used in the treatment of diarrhoea, asthma and passing of blood in urine; the bark is used as poultice in muscular pain; the roots, barks and leaves are used in epilepsy (Kirtikar and Basu, 1980; Uddin *et al.*, 2008). In African folk medicine, it is used in many diseases

*Correspondence: Sarder Nasir Uddin, Department of Biology and Chemistry, Graduate School of Medicine, Yamaguchi University (Yoshida Campus), 1677-1 Yoshida Yamaguchi-735-8512, Japan. E-mail: s_n_uddin@yahoo.co.jp

including dysentery, hypertension, etc (Iwe, 1993) Fruit, leaves, bark, stems, twigs and seeds are also used in traditional medicine. The leaves are used to treat coughs and sore throats and the bark is used to make cough syrups. Other reported uses include remedies for bronchitis, gonorrhoea, malaria, yellow fever, toothaches, and intestinal worms (Rulangaranga, 1991; Uddin *et al.*, 2008).

Free radicals are metastable chemical species which, after being generated *in vivo* as by products of various biochemical reactions, tend to rob electrons from the molecules in the immediate surrounding in order to replace their own losses. These radicals may be envisaged as molecular sharks, which if not scavenged effectively on time, are capable of damaging crucial bio-molecules including those present in cell membranes, mitochondria, DNA etc. and thus predisposing various pathophysiological states. The role of free radicals, especially of the so called 'reactive oxygen species' (ROS), has been well-established in the pathogenesis of many disease conditions such as rheumatoid arthritis, hemorrhagic shock, cardiovascular disorders, cystic fibrosis, some metabolic disorders, neurodegenerative diseases (e.g. Parkinsonism, Alzheimer's disease), gastrointestinal ulcerogenesis, AIDS and even early senescence (Halliwell and Gutteridge, 1985; Halliwell, 1994). ROS is a collective term, which includes not only the oxygen radicals (O_2^- and OH) but also some non-radical derivatives of oxygen. These include hydrogen peroxide (H_2O_2), hypochlorous acid (HOCl) and ozone (O_3). In recent years one of the areas which attracted a great deal of attention is the possible therapeutic potential of antioxidants in controlling degenerative diseases associated with marked oxidative damage. Several plant extracts and different classes of phytochemicals have been found to have quite prominent antioxidant activity (Larson, 1988; Tripathi *et al.*, 1996; Sreejayan and Rao, 1997; Vani *et al.*, 1997). Methanol extract of *Senna tora* possess strong antioxidant activity (Uddin *et al.*, 2008a) The objective of the present study was to investigate the

antioxidant and antimicrobial activity of the crude extract of *Trema orientalis*.

During the past decade, traditional systems of medicine have become increasingly important in view of their safety. Current estimates suggest that, in many developing countries, a large proportion of the population relies heavily on traditional practitioners and medicinal plants to meet primary health care needs. Although modern medicine may be available in these countries, herbal medicines (phytomedicines) have often maintained popularity for historical and cultural reasons. Concurrently, many people in developed countries have begun to turn to alternative or complementary therapies, including medicinal herbs (Farnsworth and Soejarto, 1991).

MATERIALS AND METHODS

Plant materials

Fresh leaves of *T. orientalis* were collected from Khulna University Campus in Bangladesh. The plant was identified by the experts of Bangladesh National Herbarium, Mirpur, Dhaka (Accession no. 31,285) and a voucher specimen was also deposited there. The fresh leaves were cleaned, dried and pulverized. About 400 g of powdered material was taken in a clean, flat bottomed glass container (4 L) and soaked in 1300 ml of 80% of methanol. The container was sealed and kept for a period of 10 days with occasional shaking and stirring. Then it was filtered and concentrated by evaporation.

Determination of antioxidant activities

The anti-oxidant potential of the ethanolic extract was determined on the basis of their scavenging activity of the stable 1,1-diphenyl-2-picryl hydrazyl (DPPH) free radical. DPPH is a stable free radical containing an odd electron in its structure and usually utilized for detection of the radical scavenging activity in chemical analysis. The aliquots of the different concentrations (1 - 500 μ g/ml) of the extract

were added to 3 ml of a 0.004% w/v solution of DPPH. Absorbance at 517 nm was determined after 30 min, and IC₅₀ (Inhibitory concentration 50%) was determined. IC₅₀ value denotes the concentration of sample required to scavenge 50% of the DPPH free radicals (Gupta *et al.*, 2003).

At first 6 test tubes were taken to make aliquots of 6 conc. (1, 5, 10, 50, 100 and 500 µg/ml). Plant extract and ascorbic acid were weighed 3 times and dissolved in ethanol to make the required concentration by dilution technique. Here ascorbic acid was taken as standard. DPPH was weighed and dissolved in ethanol to make 0.004% (w/v) solution. To dissolve homogeneously magnetic stirrer was used. After making the desired concentrations 3 ml of 0.004% DPPH solution was applied on each test tube by pipette. The room temperature was recorded and kept the test tubes for 30 min in light to complete the reactions. DPPH was also applied on the blank test tubes at the same time where only ethanol was taken as blank. After 30 min, absorbance of each test tubes were determined by UV spectrophotometer. IC₅₀ was determined from % inhibition vs concentration graph.

Determination antibacterial activities

Nutrient agar media was prepared by adding water to a dehydrated product that contains all the ingredients. Practically all media are available commercially in powdered form (Pelczar *et al.*, 1993).

Three types of discs were prepared for antibacterial screening: One gram sample extracts was dissolved in 10 ml of ethanol to prepare sample solution, 0.03 gm/10 ml gentamycin standard disc used as positive control to ensure the activity of standard antibiotic against the test organisms as well as for comparison of the response produced by known antibacterial agent with that produced by test samples, and third one was a blank sample (only ethanol) which was used as negative control to ensure that the residual solvents was not active. Specific organisms were inoculated into previously

sterilized nutrient agar media, mixed thoroughly and transferred immediately to the sterile petri dish in an aseptic condition. It was stored in an incubator for about 24 h to allow the proper growth of microbes. Prepared sample solutions were applied to the corresponding cups or holes with the help of a micropipette. The plates were then allowed to stand to diffuse the sample solution into the antibiotic medium at room temperature for 2 h. The plates were then incubated at 37 °C for overnight. After proper incubation, clear zones of inhibition around the point of application of sample solution were formed. These inhibition zones were measured by slide calipers and expressed in millimeter (Bauer *et al.*, 1966).

RESULTS AND DISCUSSION

For Antioxidant Activities test

DPPH is the best, easiest and widely used method for testing preliminary free radical scavenging activity of a compound or a plant extract. In the present study, methanol extracts of the leaves of *T. orientalis* showed potential free-radical scavenging activity but aqueous extract showed very little free-radical scavenging activity. The free radical scavenging property may be one of the mechanisms by which this drug is effective as a traditional medicine. Most of the tannins and flavonoids are phenolic compounds and may be responsible for antioxidant properties of many plants (Larson, 1988). IC₅₀ of the methanol extract of *T. orientalis* was 110.25 µg/ml which indicated the strong antioxidant activity of the plant extract. However the aqueous extract showed mild antioxidant activity (Figs. 1 & 2).

For Antibacterial Activities Test

Crude methanol extract of *Trema orientalis* showed antibacterial activity against *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Plesiomonas shigelloides*, *Shigella dysenteriae*, and *Vibrio cholerae* on the other hand aqueous extract showed antibacterial activity against *Staphylococcus aureus*, *Staphylococcus epidermidis*,




Fig. 1. DPPH Scavenging Assay of *T. orientalis* compared with Standard ascorbic acid (absorbance vs concentration).




Fig. 2. DPPH Scavenging Assay of *T. orientalis* compared with standard ascorbic acid (% of inhibition vs concentration).

Staphylococcus saprophyticus, *Streptococcus pyogenes* *Plesiomonas shigelloides*, *Shigella dysenteriae*, *Vibrio*

cholerae, *Shigella Flexner*, *Shigella sonnei* and *Pseudomonas aeruginosa*. Both extracts did not show any activities against *Salmonella typhi* and *Shigella boydii*.

The result of the antibacterial activity measured in term of diameter of zone of inhibition in mm. Standard antibiotic discs of Gentamycin was used as standard comparison purpose. Both extract showed antibacterial activity against both gram positive and gram negative bacteria. Aqueous extract showed higher anti microbial activities than methanol extract. The reputation of *T. orientalis* as a remedy for different microbial diseases traditionally including diarrhoea and dysentery was supported by the antibacterial screening tests. Inference can be drawn that the antibacterial constituents are present in the extract in moderate concentration (Table 1).

CONCLUSION

Methanol extract of *T. orientalis* possess strong antioxidant activity. However the aqueous extract showed mild antioxidant activity. This activity may be due to the presence of phenolic compounds

Table 1. Antibacterial activity of methanol and aqueous extract of leaves of Trema orientalis

Name of bacteria	Diameter zone of inhibition in mm		
	Gentamycin (30 μg/well)	Methanol extract (500 μg/well)	Aqueous extract (500 μg/well)
Gram positive bacteria			
<i>Staphylococcus aureus</i>	23	9	11
<i>Staphylococcus epidermidis</i>	21	9	11
<i>Staphylococcus saprophyticus</i>	32	-	13
<i>Streptococcus pyogenes</i>	21	-	10
Gram negative bacteria			
<i>Plesiomonas shigelloides</i>	24	8	9
<i>Shigella dysenteriae</i>	24	9	10
<i>Vibrio cholerae</i>	28	9	9
<i>Salmonella typh</i>	31	-	-
<i>Shigella flexneri</i>	21	-	10
<i>Shigella boydii</i>	23	-	-
<i>Shigella sonnei</i>	24	-	9
<i>Pseudomonas aeruginosa</i>	27	-	10

'-' No inhibition

(tannins and flavonoids) present in the extract. (Sadhu *et al.*, 2003). Both methanol and aqueous extract of *T. orientalis* show significant antibacterial activity against few gram positive and gram negative bacterial strains. Bangladesh imports a large quantity of pharmaceutical raw materials including medicinal plants and semi processed plant products to produce drugs and medicines. This huge foreign exchange can be saved if the indigenous medicinal plants or their semi-processed products are utilized by the manufacturers to satisfy their needs. So, further pharmacological and toxicological study is required to establish the therapeutic uses of the plant and particularly with its active principles.

REFERENCES

- Bauer AW, Kirby WMM, Sherris JC, Turck M. (1966) Antibiotic susceptibility testing by a standardized single disc method. *Am. J. Clin. Pathol.* **45**, 493-496.
- Farnsworth NR, Soejarto DD. (1991) Global importance of medicinal plants. In: Akerele O, Heywood V, Synge H. (Eds.) "The Conservation of Medicinal Plants", Cambridge University Press, Cambridge, 25-51.
- Gupta M, Mazumdar UK, Sivakumar T, Vamis MLM, Karkis S, Sambathkumar R, Mainkandan L. (2003) Antioxidant and Anti-inflammatory activities of *Acalypha Fruticosa*. *Nig. J. Nat. Prod. Med.* 25-29.
- Halliwell H. (1994) Free radicals, antioxidants, and human disease: Curiosity, cause or consequence? *Lancet* **344**, 721-724.
- Halliwell B, Gutteridge JMC. (1985) Free radicals. In Biology and Medicine, 2nd ed., Clarendon Press, Oxford, p.12-109.
- Iwe MM. (1993) Hand Book of Medicinal Plants. CRC Publication. p. 251.
- Kirtikar KR, Basu BD. (1980) Indian Medicinal Plants. 2nd Ed, Vol. 1, p. 264.
- Larson RA. (1988) The antioxidants of higher plants. *Phytochemistry* **27**, 969-978.
- Pelczar MJ, Chan JR, Noel ECS, Krieg KR. (1993) Microbiology. Fifth edition. Tata McGraw-Hill Publishing Company Ltd., New Delhi, India. pp.103-105, 510-539.
- Rulangaranga ZK. (1991) Conservation of medicinal and aromatic plants in Tanzania proceedings of a workshop on priority species for tree planting and afforestation in Tanzania, Morogoro, Tanzania. May 14-18.
- Sreejayan Rao MNA. (1997) Nitric oxide scavenging by curcuminoids. *J. Pharm. Pharmacol.* **49**, 105-107.
- Tripathi YB, Chaurasia S, Tripathi E, Upadhyay A, Dubey GP. (1996) *Bacopa monniera* Linn. As an antioxidant: Mechanism of action. *Indian J. Exptl. Biol.* **34**, 523-526.
- Uddin SN, Uddin KMA, Ahmed F. (2008) Analgesic and antidiarrhoeal activities of *Trema orientalis* Linn. in mice, *Orient. Pharm. Exp. Med.* **8**, 187-191.
- Uddin SN, Ali ME, Yesmin MN. (2008a) Antioxidant and Antibacterial Activities of *Senna tora* Roxb, *Am. J. Plant Physiol.* **3**, 96-100.
- Vani T, Rajani M, Sarkar S, Shishoo CJ. (1997) Antioxidant properties of the ayurvedic formulation Triphala and its constituents. *Int. J. Pharmacog.* **35**, 313-317.